

# Journal of Skin and Sexually **Transmitted Diseases**

Article in Press



Net Image

# Ultraviolet-induced fluorescence trichoscopy of trichobacteriosis axillaris

Vishal Gaurav<sup>1</sup>, Vaishnavi Modi<sup>1</sup>, Deepika Yadav<sup>1</sup>

<sup>1</sup>Department of Dermatology and Venereology, Maulana Azad Medical College, New Delhi, India.

#### \*Corresponding author:

Deepika Yadav, Department of Dermatology and Venereology, Maulana Azad Medical College, New Delhi, India.

deepikayadav18.90@gmail.com

Received: 20 June 2024 Accepted: 21 July 2024 EPub Ahead of Print: 09 August 2024 Published:

#### DOI

10.25259/JSSTD\_21\_2024

**Quick Response Code:** 



A 23-year-old female presented with complaint of malodour from bilateral axillae for two years, which worsened during summers. Examination revealed firmly adherent, yellow-white concretions along multiple hair shafts in both axillae associated with an offensive smell [Figure 1]. These concretions showed bluish fluorescence under Wood's lamp [Figure 2]. Polarized trichoscopy showed pale-yellow, cotton-like waxy structures forming sheaths, nodules, and concretions along the hair shafts [Figure 3], which showed bluish fluorescence on ultraviolet-induced fluorescence trichoscopy [Figure 4]. Microscopic examination of 10% potassium hydroxide mount revealed irregular cottony



Figure 1: Firmly adherent, yellow-white concretions along multiple hair shafts over right axilla.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms. ©2024 Published by Scientific Scholar on behalf of Journal of Skin and Sexually Transmitted Diseases



Figure 2: Bluish fluorescence under Wood's lamp.



Figure 3: Polarized trichoscopy showing pale-yellow, cotton-like waxy structures forming sheaths, nodules, and concretions along the hair shafts (DermLite DL5, ×20).

concretions along hair shafts. She was diagnosed with trichobacteriosis axillaris (trichomycosis axillaris) based on the above findings. The patient was advised to shave her axillary hair and prescribed clindamycin phosphate 1% gel for twice daily topical application resulting in remission within two weeks.



Figure 4: Ultraviolet-induced fluorescence trichoscopy showing bluish fluorescent concretions (DermLite DL5, ×20).

#### **Ethical approval**

The Institutional Review Board approval is not required.

## Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent.

## Financial support and sponsorship

#### **Conflicts of interest**

There are no conflicts of interest.

# Use of artificial intelligence (AI)-assisted technology for manuscript preparation

The authors confirm that there was no use of artificial intelligence (AI)-assisted technology for assisting in the writing or editing of the manuscript and no images were manipulated using AI.

How to cite this article: Gaurav V, Modi V, Yadav D. Ultraviolet-induced fluorescence trichoscopy of trichobacteriosis axillaris. J Skin Sex Transm Dis. doi: 10.25259/JSSTD\_21\_2024