



Symposium

# Topical and intralesional immunotherapy in cutaneous infections

Bini Chandran

Department of Dermatology, Ahalia Hospital, Musaffah, Abu Dhabi, United Arab Emirates.

**\*Corresponding author:**

Bini Chandran,  
Department of Dermatology,  
Ahalia Hospital, Musaffah, Abu  
Dhabi, United Arab Emirates.  
drbinichandran@gmail.com

Received: 24 April 2021  
Accepted: 14 May 2021  
Epub Ahead of Print: 10 January 2022  
Published: 09 April 2022

DOI  
10.25259/JSSTD\_36\_2021

Quick Response Code:



## ABSTRACT

Immunotherapy has revolutionized the treatment of extensive and resistant warts. Promising results have extended the role of immunotherapy to other infections such as extensive molluscum contagiosum, recurrent herpes simplex infections, and cutaneous leishmaniasis, which are resistant to standard treatment. This review focuses on topical and intralesional immunotherapy in the management of cutaneous infections.

**Keywords:** Immunotherapy, Topical, Intralesional, Wart, Molluscum contagiosum

## INTRODUCTION

Agents used for immunotherapy act by enhancing or suppressing the immune system.<sup>[1]</sup> Immune enhancing or immune-modulating agents are found to be effective in the management of recalcitrant warts.<sup>[1]</sup> Molluscum contagiosum, recurrent herpes labialis, and genitalis and cutaneous leishmaniasis are the other cutaneous infections where immunotherapy is found effective.<sup>[2-10]</sup>

This review focuses on the topical/intralesional agents used in cutaneous infections. The common topical agents used are imiquimod 5% cream, diphenylcyclopropanone (DPCP) (diphencyprone), squaric acid dibutyl ester (SADBE), 15% sinecatechins ointment, Bacillus Calmette–Guerin (BCG) vaccine in normal saline or salicylic acid, and 15% zinc ointment.<sup>[1,11-13]</sup> Intralesional agents found effective are *Mycobacterium indicus pranii* vaccine, BCG vaccine, tuberculin purified protein derivative (PPD), measles-mumps-rubella (MMR) vaccine, Candida extract, Trichophyton antigen, vitamin D3, zinc sulfate, and interferon alfa (IFN  $\alpha$ )-2b.<sup>[1,14]</sup> Autoinoculation (autoimplantation) is another promising therapy that has achieved clearance of warts by stimulating the immune response.<sup>[1,14]</sup>

### Mechanism of action: An overview

Imiquimod, an agonist of toll-like receptor 7, exerts antiviral effects by increasing the cellular levels of IFN- $\alpha$ , tumor necrosis factor-alpha, and interleukin-6.<sup>[1,15]</sup> The mechanism of action of SADBE and diphencyprone (DPCP) is not fully explained. The possible mechanisms include a type IV hypersensitivity reaction or a non-specific inflammatory reaction.<sup>[16]</sup> Cytokines released during the allergic reaction caused by the contact sensitizers tend to activate the natural killer cells which in turn lyse the virus-infected cells.<sup>[17]</sup>

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.  
©2022 Published by Scientific Scholar on behalf of Journal of Skin and Sexually Transmitted Diseases

The therapeutic effects of sinecatechins in cutaneous infections are attributed to their immunomodulatory, antioxidant, and antiviral properties.<sup>[2]</sup> The beneficial effect of zinc is believed to be due to its effects on macrophage and neutrophil functions, natural killer cell/phagocytic activity, and various cytokines.<sup>[13]</sup>

The various vaccines act by boosting cell-mediated immunity, while IFN  $\alpha$ -2b shows antiviral activity through viral interference. In addition, it has immunomodulatory and antiproliferative properties.<sup>[1]</sup>

## IMMUNOTHERAPY FOR WARTS

Recalcitrant, recurrent, or extensive warts as well as warts affecting periungual and palmoplantar sites (difficult to treat areas) are considered for immunotherapy.<sup>[1]</sup> The first contact allergen that was used in the treatment of resistant warts was dinitrochlorobenzene (DNCB) in 1973.<sup>[18]</sup> DNCB is no longer used widely due to its mutagenic potential.<sup>[11]</sup>

Autoimplantation (harvesting of wart tissue followed by autoimplantation) and Falknors needling method (index wart is punctured several times with a 26-gauge needle to produce a “beefy” red wound) are the other methods that act by modulating the immune system.<sup>[19]</sup>

The various topical and intralesional immunotherapy agents used in warts and the dose and frequency of administration are shown in Table 1.<sup>[1,12,13,16-18,20,21]</sup>

Among topical agents, an evidence-based review done by Ahn and Huang noted that clinical improvement was seen in 33–50% of patients who received 5% imiquimod cream, although none of the patients experienced complete clinical clearance.<sup>[22]</sup> A quantitative systematic review on genital warts reported that imiquimod achieved complete clearance in 51% of patients, but was not effective in human immunodeficiency virus (HIV) positive patients.<sup>[23]</sup>

Among immune sensitizers, SADBE is considered the safest – with complete clearance in 58% of patients.<sup>[24]</sup>

Sinacatechins are FDA approved for genital warts.<sup>[25]</sup> It is prepared from green tea leaves from the plant *Camellia sinensis*. A blend of eight different catechins and other green tea components constitute the ointment. Epigallocatechin-3-gallate is the main catechin in the preparation.<sup>[26]</sup> The clearance rate of genital warts documented following application of 15% sinacatechins ointment varied from 46% to 52% and recurrence rate was low at 6.5%.<sup>[27]</sup>

Topically applied BCG was effective in children with common warts and plane warts with 65% and 45% resolution, respectively.<sup>[28]</sup> In a placebo-controlled study, 80% of the patients had complete resolution of genital warts after a maximum of six BCG applications.<sup>[29]</sup>

Songsantiphap and Asawanonda, after assessing the efficacy of 15% zinc oxide ointment in 16 patients, found that the ointment applied three times a day for 4 weeks attained reduction in size of warts in comparison to placebo. The authors concluded that zinc oxide ointment can serve as adjunctive therapy to increase the efficacy of other modalities of treatment.<sup>[13]</sup>

A case series on recalcitrant cutaneous warts treated with intralesional MMR vaccine reported complete response in nine of the eleven treated (81.9%), which was comparable to the 82% complete clearance reported by Chauhan *et al.*<sup>[30,31]</sup> In an open, uncontrolled trial with intralesional vitamin D3 therapy in cutaneous warts, 78.5% of patients had complete resolution.<sup>[32]</sup> In a 2 year study of cutaneous warts treated with intralesional Candida antigen, 80% of patients showed complete clearance of lesions. Six out of the seven immunocompromised patients who were treated also demonstrated either a partial or complete response.<sup>[33]</sup> Hypersensitivity to Candida antigen is an absolute contraindication for this treatment modality.<sup>[1]</sup> In a randomized controlled study, 62% of the patients responded to Trichophyton when used as a single agent.<sup>[34]</sup> *Mycobacterium w* vaccine was found to have comparable efficacy to cryotherapy with 66.7% clearance in a study by Dhakar *et al.*<sup>[35]</sup> Sharquie *et al.* demonstrated 39.7% clearance of cutaneous warts in a single-blinded placebo-controlled trial using BCG vaccine.<sup>[36]</sup> Podder *et al.* in a double-blind, randomized controlled trial that assessed the effectiveness and safety of intradermal BCG vaccine in comparison to intradermal tuberculin PPD, observed both agents to be useful with better response in the BCG treated group (complete clearance in 48.5% of the BCG treated group and in 18.5% of the tuberculin PPD treated group).<sup>[37]</sup>

A randomized controlled trial reported the usefulness of Propionium bacterium parvum (0.1 ml injected intralesionally once a month for 3–5 months) in cutaneous warts.<sup>[21]</sup>

Another method attempted for warts is autoimplantation. In autoimplantation, a full-depth nick up to subcutis level (under local anesthesia and aseptic precautions) is made with an 18-gauge needle in a selected wart which serves as the donor wart. A chunk of wart tissue is removed and introduced in a subcutaneous pocket created with the same 18-gauge needle on the flexor aspect of the left forearm about 2 inches below the antecubital crease and secured in place by band-aid plaster. The dressings are removed after 5 days.<sup>[38]</sup> Autoimplantation achieved clearance of lesions in 91% of patients at 2 months.<sup>[38]</sup> ElGhareeb compared autoimplantation with MMR vaccine and noted that autoimplantation showed better efficacy in achieving clearance of distant warts (warts that were situated away from the treated wart).<sup>[39]</sup> Complete resolution was noted in 70.7% patients with palmoplantar warts treated with Falknors needling method.<sup>[19]</sup>

**Table 1:** Topical and intralesional immunotherapy agents used in warts.

Route of administration	Drug used	Dose, frequency of administration, and duration of treatment	Indication
Topical agents	Imiquimod	5% cream three times a week for 16 weeks	Cutaneous and genital warts
	Sinecatechins	15% ointment – three times a week for 16 weeks	Cutaneous warts
	Bacillus Calmette-Guerin	Application in normal saline or salicylic acid, wash off after 2 hours, weekly treatment for up to 6 to 12 weeks	Cutaneous and genital warts
	Topical zinc	15% ointment three times a day for 4 weeks	Cutaneous warts
	Topical interferon	Human leukocyte $\alpha$ -interferon (incorporated $2 \times 10^6$ IU/g) in hydrophilic cream, 6 gram thrice a day for 4 consecutive days a week up to 4 weeks	Vulvar and vaginal warts
	Squaric acid dibutyl ester (SADBE)	Paring of warts followed by application of SADBE. Occlude with dressings for four hours and then wash with soap and water. Repeat sessions every 2–4 weeks or till resolution	Cutaneous warts
	Diphencyprone	Sensitize with 2% and follow-up by weekly maintenance of 0.001–1% diphencyprone. Occlude with dressings for 48 hours and wash with soap and water. Repeat sessions at intervals of 2–4 weeks or till resolution	Cutaneous warts
Intralesional agents	Measles-mumps-rubella vaccine	0.3–0.5 ml into the largest wart, repeat every 2 weeks for 3 to 4 sessions	Cutaneous warts
	Vitamin D	0.2 ml of 7.5 mg/ml, 2 sittings 4 weeks apart	Cutaneous and genital warts
	Candida antigen	0.1–0.3ml into the largest wart, then 3 sessions at weekly intervals	Cutaneous warts
	Trichophyton antigen	0.3 ml into the largest wart every 3 weeks, maximum 5 sittings	Cutaneous and genital warts
	Mycobacterium Indicus Pranii vaccine	0.1 ml into 3–5 warts or all warts followed by up to 10 sessions 2–4 weekly	Cutaneous warts
	Bacillus Calmette-Guerin vaccine	0.1–0.5 ml into the largest wart, every 2 weeks for 5 sittings	Cutaneous and genital warts
	Tuberculin purified protein derivative	2.5 units into a few warts every 2 weeks	Cutaneous warts
	Interferon alfa-2b	1–2 million units 3 days/week for 3 weeks	Genital warts
	Propionium bacterium parvum	0.1 ml once a month for 3–5 months	Cutaneous warts

A network meta-analysis on intralesional immunotherapy for warts concluded that PPD and MMR showed superior efficacy in achieving complete primary and distant recovery and showed reduced recurrence rate at the same site in comparison to cryotherapy and other immunotherapeutic modalities.<sup>[14]</sup> Autoinoculation showed efficacy comparable to MMR and PPD in attaining distant recovery.<sup>[14]</sup>

Unlike PPD, intralesional MMR vaccine is not recommended in pregnancy.<sup>[40]</sup>

### Molluscum contagiosum

Being a self-limiting condition, opinions differ on whether to treat molluscum contagiosum or not. According to general consensus, patients with extensive disease (mainly immunosuppressed individuals), secondary bacterial infection, conjunctivitis, molluscum dermatitis, or esthetic complications require treatment.<sup>[41]</sup>

The topical immunomodulatory methods that have been tried in molluscum contagiosum are 5% imiquimod cream, diphencyprone, and topical sinecatechins.<sup>[2-7]</sup> Al-Mutairi *et al.*, in a prospective, randomized, comparative, and observer-blinded study, found imiquimod 5% cream applied 5 days a week to be of comparable efficacy (complete cure rate 91.8%) to weekly cryotherapy (complete cure rate 100%) at the end of 16 weeks. Pain, bullae formation, pigmentary changes, and superficial scarring were significantly more in the cryotherapy group. The authors concluded that imiquimod could be the preferred option, especially in children with several small lesions.<sup>[3]</sup> However, van der Wouden *et al.* found imiquimod to be not more effective than placebo.<sup>[42]</sup>

Kang *et al.* studied the effect of diphencyprone solution applied weekly starting with 0.0001% and increased to a maximum of 0.1%, based on the response observed. The authors documented complete cure in 14/22 children (63.6%) within a mean treatment period of 5.1 weeks. Although most

**Table 2:** Intralesional immunotherapy agents used in molluscum contagiosum.

Intralesional immunotherapy agent	Dose and frequency	Percentage of patients who achieved complete resolution	Serious side effects
Measles-mumps-rubella vaccine	0.3 ml into the largest lesion at an interval of 2 weeks	Two patients treated - complete resolution in both (100%) after two and four injections, respectively	None
Candida antigen	0.3 ml into 1–3 lesions, at an interval of 2–6 weeks	50–56%	None
Purified protein derivative	10 IU (0.1 ml) into the largest lesion every 2 weeks up to six injections	85%	One case of anaphylaxis in 20 patients treated
Vitamin D	0.2ml into two relatively large lesions at 4 weeks interval	Complete resolution of all lesions in two immunosuppressed patients (100%) after 2 injections	None

of the treated children could tolerate the treatment (most common adverse effects being erythema and pruritus), four patients had to discontinue diphencyprone.<sup>[4]</sup>

Sinecatechins 10% ointment twice daily for 4 weeks applied to the top of the lesions was found effective in a 5-year-old child with extensive molluscum lesions.<sup>[2]</sup>

Intralesional immunotherapies [Table 2] used in molluscum contagiosum include measles-mumps-rubella vaccine, Candida, tuberculin PPD, vitamin D3, interferon  $\alpha$ , and Streptococcal substrain OK-432.<sup>[5-7]</sup>

Nelson *et al.* reported 10 patients with HIV infection who received intralesional IFN  $\alpha$  (one mega unit) at weekly intervals for 4 weeks. Three Mollusca of each patient were chosen for treatment. Complete resolution was seen in 36.7% of the treated lesions. A reduction in size (50–90%) was observed in 18 lesions (18/30, 60%), while one lesion remained unchanged (3.3%).<sup>[5]</sup> Inui *et al.* reported successful treatment of molluscum contagiosum with intralesional penicillin-treated and heat-treated lyophilized powder of a substrain of *Streptococcus pyogenes* A (OK 432) in a patient receiving chemotherapy.<sup>[6]</sup>

After reviewing 10 studies on intralesional immunotherapy for molluscum contagiosum, Wells *et al.* concluded that none of the patients who completely responded to Candida, MMR vaccine, OK-432, and vitamin D3 showed recurrence after immunotherapy, while intralesional PPD and interferon  $\alpha$  showed less efficacy in preventing recurrences.<sup>[7]</sup>

## RECURRENT HERPES SIMPLEX INFECTION

A double-blind, randomized placebo-controlled study in patients who had six or more self-reported episodes of herpes labialis in the preceding 12 months suggested that topical sensitization dose of SADBE (2%) on the arm may prevent herpes simplex virus (HSV) outbreaks and this was supported by another multicentre study.<sup>[8,9]</sup>

Bernstein *et al.* after evaluation of subunit vaccines and imiquimod in guinea pig model of recurrent HSV type-2 infection concluded that the strategy of prime (subunit HSV

vaccine) and topical pull (intravaginal/topical imiquimod) decreased recurrent HSV more effectively than vaccine alone.<sup>[43]</sup>

## CUTANEOUS LEISHMANIASIS

Nahidi *et al.* found a combination of topical diphencyprone and intralesional meglumine antimoniate to be more effective in acute urban cutaneous leishmaniasis than meglumine antimoniate administered alone.<sup>[44]</sup> Topical imiquimod (5–7.5%) in combination with meglumine antimoniate was able to cure cutaneous leishmaniasis in patients who did not respond to meglumine antimoniate alone.<sup>[10,15,45]</sup>

## CONCLUSION

With the emergence of pathogens resistant to commonly used antimicrobials, immunotherapy alone or in combination with other modalities appears to be the need of the hour. Immunotherapy could be the key to treatment-resistant infections and focused research in this area may uncover lasting solutions.

## Declaration of patient consent

Not required as there are no patients in this article.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Thappa DM, Chiramel MJ. Evolving role of immunotherapy in the treatment of refractory warts. *Indian Dermatol Online J* 2016;7:364-70.
- Padilla Espana L, Mota-Burgose A, Martinez-Amo JL, Benevente-Ortiz F, Rodriguez-Bujaldon A. Recalcitrant

- molluscum contagiosum successfully treated with sinecatechins. *Dermatol Ther* 2016;29:217-8.
3. Al-Mutairi N, Al-Doukhi A, Al-Farag S, Al-Haddad A. Comparative study on the efficacy, safety, and acceptability of imiquimod 5% cream versus cryotherapy for molluscum contagiosum in children. *Pediatr Dermatol* 2010;27:388-94.
  4. Kang SH, Lee D, Park J, Cho SH, Lee SS, Park SW. Treatment of molluscum contagiosum with topical diphencyprone therapy. *Acta Derm Venereol* 2005;85:529-30.
  5. Nelson MR, Chard S, Barton SE. Intralesional interferon for the treatment of recalcitrant molluscum contagiosum in HIV antibody positive individuals-a preliminary report. *Int J STD AIDS* 1995;6:351-2.
  6. Inui S, Asada H, Yoshikawa K. Successful treatment of molluscum contagiosum in the immunosuppressed adult with topical injection of streptococcal preparation OK-432. *J Dermatol* 1996;23:628-30.
  7. Wells A, Saikaly SK, Schoch JJ. Intralesional immunotherapy for molluscum contagiosum: A review. *Dermatol Ther* 2020;33:e14386.
  8. Palli MA, McTavish H, Kimball A, Horn TD. Immunotherapy of recurrent herpes labialis with squaric acid. *JAMA Dermatol* 2017;153:828-9.
  9. Lynn AS, Honari G, Guan L, Zhao L, Palli MA, Horn TD, *et al.* A phase 2, multi-center, placebo-controlled study of single dose squaric acid dibutyl ester (SADBE) to reduce frequency of outbreaks in subjects with recurrent herpes labialis. *J Am Acad Dermatol* 2020;83:1807-9.
  10. Azim M, Khan SA, Ullah S, Ullah S, Anjum SI. Therapeutic advances in the topical treatment of cutaneous leishmaniasis: A review. *PLoS Negl Trop Dis* 2021;15:e0009099.
  11. Suh DW, Lew BL, Sim WY. Investigations of the efficacy of diphenylcyclopropanone immunotherapy for the treatment of warts. *Int J Dermatol* 2014;53:e567-71.
  12. Word AP, Nezafati KA, Cruz PD Jr. Treatment of warts with contact allergens. *Dermatitis* 2015;26:32-7.
  13. Songsantiphap C, Asawanonda P. Topical 15% zinc oxide ointment significantly reduces the size of common warts after four weeks: A randomized triple blinded placebo controlled study. *J Clin Aesthet Dermatol* 2019;12:26-31.
  14. Salman S, Ahmed MS, Ibrahim AM, Mattar OM, El-Shirbiny H, Sarsik S, *et al.* Intralesional immunotherapy for the treatment of warts: A network meta-analysis. *J Am Acad Dermatol* 2019;80:922-30.
  15. Miranda-Verastegui C, Tulliano G, Gyorkos TW, Calderon W, Rahme E, Ward B, *et al.* First-line therapy for human cutaneous leishmaniasis in Peru using the TLR7 agonist imiquimod in combination with pentavalent antimony. *PLoS Negl Trop Dis* 2009;3:e491.
  16. Kehdi J, Erickson C, Rady P, Tyring S, Gaspari AA. Epidermodysplasia verruciformis: Successful treatment with squaric acid dibutyl ester. *Cutis* 2015;96:114-8.
  17. Pollock B, Highet AS. An interesting response to diphencyprone (DPC) sensitization on facial warts: Review of DPC treatment for viral warts. *J Dermatol Treat* 2002;13:47-50.
  18. Lewis HM. Topical immunotherapy of refractory warts. *Cutis* 1973;12:863-7.
  19. Kumari P, Yadav D, Vijay A, Jain SK, Kumar M, Kumar R, *et al.* Falknor's needling method as a potential immunotherapy in palmo-plantar warts. *Indian J Dermatol Venereol Leprol* 2019;85:129.
  20. Syed TA, Lundin S, Cheema KM, Kahlon BM, Cheema R, Ahmad SA. Human leukocyte interferon- $\alpha$  in cream, for the treatment of genital warts in Asian women a placebo-controlled, double-blind study. *Clin Investig* 1994;72:870-3.
  21. Nasser N. Treatment of common warts with the immune stimulant propionium bacterium parvum. *An Bras Dermatol* 2012;87:5859.
  22. Ahn CS, Huang WW. Imiquimod in the treatment of cutaneous warts: An evidence-based review. *Am J Clin Dermatol* 2014;15:387-99.
  23. Moore RA, Edwards JE, Hopwood J, Hicks D. Imiquimod for the treatment of genital warts: A quantitative systematic review. *BMC Infect Dis* 2001;1:3.
  24. Silverberg NB, Lim JK, Paller AS, Mancini AJ. Squaric acid immunotherapy for warts in children. *J Am Acad Dermatol* 2000;42:803-8.
  25. Fantasia HC. Sinecatechins ointment 15% for the treatment of external genital warts. *Nurs Womens Health* 2012;16:418-22.
  26. Tyring S. Sinecatechins. *J Clin Aesthet Dermatol* 2012;5:19-26.
  27. El-Khalawany M, Shaaban D, Aboeldahab S. Immunotherapy of viral warts: Myth and reality. *Egypt J Dermatol Venerol* 2015;35:1-13.
  28. Salem A, Nofal A, Hosny D. Treatment of common and plane warts in children with topical viable bacillus Calmette-Guerin. *Pediatr Dermatol* 2013;30:60-3.
  29. Metawea B, El-Nashar AR, Kamel I, Kassem W, Shamloul R. Application of viable bacille Calmette-Guérin topically as a potential therapeutic modality in condylomata acuminata: A placebo-controlled study. *Urology* 2005;65:247-50.
  30. Chandran B. Intralesional measles, mumps, and rubella vaccine for the treatment of recalcitrant warts: A case series and review of literature. *J Skin Sex Transm Dis* 2021;3:56-61.
  31. Chauhan PS, Mahajan VK, Mehta KS, Rawat R, Sharma V. The efficacy and safety of intralesional immunotherapy with measles, mumps, rubella virus vaccine for the treatment of common warts in adults. *Indian Dermatol Online J* 2019;10:19-26.
  32. Kavya M, Shashikumar BM, Harish MR, Shweta BP. Safety and efficacy of intralesional Vitamin D3 in cutaneous warts: An open uncontrolled trial. *J Cutan Aesthet Surg* 2017;10:90-4.
  33. Alikhan A, Griffin JR, Newman CC. Use of Candida antigen injections for the treatment of verruca vulgaris: A two-year mayo clinic experience. *J Dermatolog Treat* 2016;27:355-8.
  34. Horn TD, Johnson SM, Helm RM, Roberson PK. Intralesional immunotherapy of warts with mumps, Candida, and trichophyton skin test antigens: A single-blinded, randomized, and controlled trial. *Arch Dermatol* 2005;141:589-94.
  35. Dhakar AK, Dogra S, Vinay K, Sarangal R, Kanwar AJ, Singh MP. Intralesional Mycobacterium w vaccine versus cryotherapy in treatment of refractory extragenital warts: A randomized, open-label, comparative study. *J Cutan Med Surg* 2016;20:123-9.
  36. Sharquie KE, Al-Rawi J, Noaimi A. Bacillus Calmette Guerin monotherapy of warts. *Saudi Med J* 2008;29:589-93.
  37. Podder I, Bhattacharya S, Mishra V, Sarkar TK, Chandra S, Sil A, *et al.* Immunotherapy in viral warts with intradermal

- Bacillus Calmette-Guerin vaccine versus intradermal tuberculin purified protein derivative: A double-blind, randomized controlled trial comparing effectiveness and safety in a tertiary care center in Eastern India. *Indian J Dermatol Venereol Leprol* 2017;83:411.
38. Shivakumar V, Okade R, Rajkumar V. Autoimplantation therapy for multiple warts. *Indian J Dermatol Venereol Leprol* 2009;75:593-5.
  39. ElGhareeb MI. Comparative study of autoimplantation therapy and intralesional injection of MMR vaccine in warts treatment. *Dermatol Ther* 2019;32:e13135.
  40. Eassa BI, Abou-Bakr AA, El-Khalawany MA. Intradermal injection of PPD as a novel approach of immunotherapy in anogenital warts in pregnant women. *Dermatol Ther* 2011;24:137-43.
  41. Leung AK, Barankin B, Hon KL. *Molluscum contagiosum*: An update. *Recent Pat Inflamm Allergy Drug Discov* 2017;11:22-31.
  42. van der Wouden JC, Menke J, Gajadin S, van der Sande R, Kruithof EJ, Sollie A, *et al*. Interventions for cutaneous molluscum contagiosum. *Cochrane Database Syst Rev* 2017;5:CD004767.
  43. Bernstein DI, Cardin RD, Bravo FJ, Awasthi S, Lu P, Pullum DA, *et al*. Successful application of prime and pull strategy for a therapeutic HSV vaccine. *NPJ Vaccines* 2019;4:33.
  44. Nahidi Y, Goyonlo VM, Layegh P, Taghavi F, Najafi MN. Immunomodulatory effects of topical diphencyprone for the treatment of acute urban cutaneous leishmaniasis. *J Dermatolog Treat* 2021;32:220-6.
  45. Miranda-Verastegui C, Llanos-Cuentas A, Arevalo I, Ward B, Matlashewski G. Randomized, double blind clinical trial of topical imiquimod 5% with parenteral meglumine antimoniate in the treatment of cutaneous leishmaniasis in Peru. *Clin Infect Dis* 2005;40:1395-403.

**How to cite this article:** Chandran B. Topical and intralesional immunotherapy in cutaneous infections. *J Skin Sex Transm Dis* 2022;4:27-32.